Melbourne Bladder Clinic

International Index of Erectile Function (IIEF – 5) Questionnaire

Circle the number of the response that best describes your own situation. Select only one response for each question.

OVER THE PAST 6 MONTHS:

1. How would you		VERY LOW	LOW	MODERATE	HIGH	VERY HIGH
rate your confidence in your ability to get and keep an erection?		1	2	3	4	5
2. When you had erections with sexual stimulations, how often were	NO SEXUAL ACTIVITY	ALMOST NEVER OR NEVER	A FEW TIMES (MUCH LESS THAN HALF	SOMETIMES (ABOUT HALF THE TIME)	MOST TIMES (MUCH MORE THAN HALF	ALMOST ALWAYS OR ALWAYS
your erections hard enough for penetration (entering your partner)?	0	1	THE TIME)	3	THE TIME) 4	5
3. During sexual intercourse, how often were you able to maintain your erection after	DID NOT ATTEMPT INTERCOUR SE	ALMOST NEVER OR NEVER	A FEW TIMES (MUCH LESS THAN HALF THE TIME)	SOMETIMES (ABOUT HALF THE TIME)	MOST TIMES (MUCH MORE THAN HALF THE TIME)	ALMOST ALWAYS OR ALWAYS
you had penetrated (entered) your partner?	0	1	2	3	4	5
4. During sexual intercourse, how difficult was it to maintain your	DID NOT ATTEMPT INTERCOUR SE	EXTREMELY DIFFICULT	VERY DIFFICULT	DIFFICULT	SLIGHTLY DIFFICULT	NOT DIFFICULT
erection to completion of intercourse?	0	1	2	3	4	5
5. When you attempted sexual intercourse, how often was it satisfactory for	DID NOT ATTEMPT INTERCOUR SE	ALMOST NEVER OR NEVER	A FEW TIMES (MUCH LESS THAN HALF THE TIME)	SOMETIMES (ABOUT HALF THE TIME)	MOST TIMES (MUCH MORE THAN HALF THE TIME)	ALMOST ALWAYS OR ALWAYS
λοης	0	1	2	3	4	5

SEVERE ED (<7), MODERATE ED (8-11), MILD to MODERATE ED (12-16), MILD ED (17-21), NO ED (>21)